


Uutta tietoa perunaseitistä

Paavo Ahvenniemi
















































10. Tubers with scab-like patches on the skin caused by *R. solani*.





















262 2 Asterix S

























Seittikokeet 1998-1999											
K-koetila											
Lajike=Kulta											
Siemen/ käsittely	Varsiluku kpl	Versol. %	Mukula- luku	Sato tn/ha	sl	Tärkkelys %	Seittirupi sl 0-4	Kasvuh. %	Vihert. %	Epämuot. %	I-luokka %
Uusi	4,6	5	13,1	35,8	100	16,5	0,1	1,4	1,9	0,0	97
Uusi P	4,7	3	13,9	36,6	102	16,4	0,0	1,5	2,0	0,0	96
Puhdas	3,3	52	11,3	35,4	99	16,3	2,0	12,3	3,9	1,7	83
Puhdas P	3,8	29	13,4	40,0	112	16,6	0,1	1,4	2,2	0,0	97
Seittirupinen	2,7	77	8,9	25,8	72	14,3	2,4	16,4	6,8	7,3	68
Seittirupinen P	3,8	45	12,8	37,5	105	16,2	2,3	5,0	2,1	0,5	91
P = siemenen upotuspeittäys											

Loppu


Sato g/kasvi




Juuret g




Varsiluku kpl




Mukulaluku kpl




Maavarsivioitukset %




Versolaikkua %:ssa varsista




Maavarret lkm



Maavarret paino



Varsipaino g



Varsi cm

